

## Application of Cognitive Behavior Therapy on the Nightmares of a Patient having Coronary Artery Disease

Sengupta, U., Singh, A. R., Ray, H. N.

Rajendra Institute of Medical Sciences (RIMS), Ranchi

Coronary artery disease (CAD) is associated with increased psychological distress and Nightmares have an impact on cardiac health as it worsens emotional regulation and increases distress. This study assesses the impact of Cognitive Behavior Therapy (CBT) on the Nightmares of a male patient having Coronary Artery Disease. This is a Single case experimental design. Anxiety, depression, stress, locus of control and personality have been studied as mediating factors in the patient using Hospital anxiety and depression scale, Perceived stress scale, Rotter's Locus of control scale and Type A/B Behavior Pattern Scale. The intervention disentangled the dream content to understand the psychological distress and its impact on cardiac outcome. Frequency of cardiac events decreased with decrease in frequency of nightmares and reduction in anxiety, depression and stress. Psychodynamic-informed CBT could successfully intervene upon the psychological distress of a middle-aged patient having Coronary Artery Disease.

**Keywords:** Nightmares, Coronary Artery Disease, Cognitive Behavior Therapy, Single Case Experimental Design

Disruptions in sleep patterns are defined as nonorganic sleep disorders. These include sleep walking, night terror, insomnia and hypersomnia. Nightmares, in which a person wake up from sleep and remembers frightening dreams that typically involve imagery related to significant threats to survival, security, or self-esteem, is one such nonorganic sleep disorder. My client, A. K., is a 45-year-old male Hindi speaker and Hindu, who experienced severe nightmares. The index patient comes out of REM sleep and is completely alert and aware of his surroundings immediately, which is in sharp contrast to the usual grogginess experienced during waking up from sleep. Most of his nocturnal dreams comprise scenarios where he and his near and dear ones are succumbing to a deadly disease or a terrible accident. It also includes his family suffering because his firm is losing money but he finds his actual circumstances to be very different

from his dreams when he wakes up. He does, however, feel anxious because while currently being awake, he is able to vividly recall the dreams. With time, this further increases his distress, which causes even greater cortisol release, thereby maintaining stability of his cardiac events. Studies have shown that CBT is effective in treating anxiety, depression, stress and health related quality of life in patients having Coronary Artery Disease (CAD) (Li, Buys, Ferguson et. al 2021). CBT has also proved to be effective in treating sleep dysfunctions like nightmares and parasomnia (Vincent, Giannouli & McQuarrie, 2023). However, no study has been done yet which addresses nightmares in patients having CAD itself using CBT approach. Hence this study is a novel addition to the already existing literature on psychotherapy in patients having CAD who also have sleep dysfunctions.

## Objectives

- To assess the anxiety, stress, depression, locus of control, type of personality of a patient having Coronary Artery Disease (CAD)
- To understand how nightmares can be related with the occurrence of cardiac events in a patient having Coronary Artery Disease (CAD)
- Intervening the psychological distress and its impact on the occurrence of nightmares and cardiac events in a patient having Coronary Artery Disease (CAD)

## Method

### Sample

Purposive sampling was used to recruit the present case from the in-patient unit of Cardiology Department, Rajendra Institute of Medical Sciences (RIMS), Ranchi.

### Tools

*Hospital Anxiety and Depression Scale (HADS)*: The HADS (Zigmond and Snaith, 1983) measures symptoms of anxiety and depression in patients currently admitted in the hospital. Its validity ranges from 0.73 to 0.77. There are 7 items measuring anxiety and 7 items measuring depression in this scale. It is a 4 point Likert scale.

*Perceived Stress Scale* (Cohen, Kamarck, Mermelstein, 1983) assessed how much the patients perceived daily life circumstances to be stressful and in what ways did such events cause the stress so that the researcher can find out the specific ways to help them cope with it. The Cronbach-Alpha Internal Consistency reliability of this scale is 0.71 to 0.91.

*Type A/B Behavior Pattern Scale*: The scale is used to measure if the participant has the characteristics of Type-A or Type-B personality and it is constructed in the Indian

context. It is presented in the form of a 5-point scale. The scale has two parts-Form A and Form B out of which Form A is used to measure Type A Personality. The reliability coefficient of Form A is .54. The validity for both the forms which have to be 0.73 (Dhar & Jain, 2001).


*Locus of Control Scale* (Rotter, 1966) is a 29-item inventory that measures the degree to which the individual interprets events as being a result of internal or external loci. This scale was used in a one year follow-up Indian study where non-pharmacological intervention was done on patients having alcohol dependence syndrome (Kumar and Thomas, 2007). The split-half reliability of this scale ranges from 0.65 to 0.79.

### The Psychopathology

In the initial session, a history related to the details of the patient's dreams was taken. History related to the patient's physiological and psychological mechanisms during the instances when he remembers the dream during his wakeful state was focused upon. Index patient dreamt of his business failing, his family members dying out of accidents and mishaps often during the night as a part of his nightmares. It was found that whenever the memories of the dreams comes in the patient, he starts hyperventilating as his breathing rate increases along with palpitations. It was reported that the index patient started having such imageries during his period of wakefulness since June, 2021 after his father's demise. Initially, he did not worry about it as he considered the nightmares to be rare incidents and expected them to decrease over time. However, when the nightmares continued to happen and the imageries came to the patient at a steady rate of about twice in a day, then he started becoming anxious about what if such mishaps happen in reality to his family and to himself. Eventually, this excessive worry affected his sleep patterns. He had difficulty falling asleep

during the night and often woke up for multiple times during sleep. During the morning, he woke up during the usual time and did not feel rested. He also reported to the clinician that he could only sleep for around three to four hours in the night without interruptions. This is because the vivid dreams during the night indicated increased cognitive activity in the patient. According to the Type A/B Behavior Pattern Scale (Dhar and Jain, 2001), the patient had a Type A personality. As a result, when he experienced anxiety, his stress level increased, which in turn decreased the potassium channel-dependent regulation of action potential firing. A decrease in K<sup>+</sup> channel inhibition actually increased the excitability of his lateral amygdala because K<sup>+</sup> channels typically have an inhibitory impact that causes the neuronal membrane to become hyperpolarized following an action potential. These intriguing findings suggest that, at least in part, long-term stress increased emotional reactivity in the patient, including fear and anxiety reactions, through localized effects on the excitability of amygdala neurons mediated by K<sup>+</sup> channel function. The efficiency with which other, fear-inhibiting regions, such as the hippocampus and medial prefrontal

cortex, regulate amygdala output would be diminished by these local effects inside the amygdala, which would probably form an overactive circuit with regard to fear and anxiety in the index patient (Ressler, 2010). In other words, during periods of extreme fear, the patient's amygdala becomes overly active, thereby reducing the effectiveness of prefrontal cortex. Hence, whenever the index patient has thoughts of impending danger to himself and his family, his ego is unable to tolerate the anxiety, as a result of which he represses such thoughts into the unconscious which therefore occurs in his dreams, resulting in nightmares. This is why the patient only had imageries of danger during his sleep but never had such worries during his wakefulness except only remembering the content of his dreams, leading to extreme stress and anxiety, which increased his chances of cardiac arrests. Eventually, due to his recurrent cardiac events, the patient became afraid of going to work and his family members became overly cautious about allowing him to take on stressful work for long, as it was found that whenever the patient was in stress for a few hours at a stretch or indulged into worry and overthinking, he had a sensation of pain in his chest.



### **The CBT Module used - Intervening Anxiety**

Following techniques were used to target reduction of anxiety in index patient –

1. Interrelationship between anxiety and increased cardiovascular event
2. Somatic Hyperawareness
3. Reducing existential fear associated with dying and passing away due to unexpected cardiac events
4. Differentiating among panic attack, normal anxious reactions
5. Using Mindfulness – Here and Now
6. Content of dreams were discussed with the patient – focusing on their relatedness with his daily life concerns

### **Intervening Depression**

In intervening depression in cardiovascular patients, the following difficulties are to be addressed –

1. Intervening negative affectivity by identifying the negative emotion and managing it,
2. Modifying social inhibition by intervening social anxiety in the index patient
3. Focusing on improving the quality of interpersonal relationship and improving social support network
4. Intervening on
  - Self-criticism and perfectionism;
  - unrealistic need for praise from nearly everyone in the vicinity;
  - unrealistic categorization of forceful and aggressive behavior
  - Criticism of oneself and other people
5. Intervening upon depressive mood state when the patient remembers the dreams during his wakeful state

6. Discussing about the perceived fatality of cardiovascular disease

### **Intervening Stress**

1. Respiratory Biofeedback –
2. EMG Biofeedback –
3. Mindfulness Based Stress reduction (MBSR) technique

### **Intervening Locus of control**

1. Switching from an external towards an internal center of control
2. Helping to regain a sense of control over the patient's outcomes of treatment
3. Using the patient's healthy locus of control to improve his health outcomes

### **Procedure**

Initially the clinician visited the In-patient department of Cardiology at RIMS, Ranchi. Thereafter, a patient having CAD was randomly recruited from the pool of patients visiting the OPD regularly who was male and in his middle ages. History about his heart disease and how he ended up having cardiac arrest recurrently was discussed. Stress was found to be the primary reason for his cardiac events along with Type A personality. Thereafter the patient was explained about how psychological distress not only causes but also exacerbates CAD and he was explained about the intervention module which targets stress, anxiety, depression and locus of control. Discussions were also made about how he is more prone to having psychological distress due to having Type A personality. Written consent was taken from the patient and he started participating in the intervention thereafter. During the intervention, it was found that his anxiety stems from recurrent nightmares which in turn is the reason for repeated cardiac events. Hence, intervention was tailored targeting nightmares along with other psychological distress in the patient. Baseline

and post-intervention assessment were done on him and his data was analysed.

### **The Treatment Process And Outcome**

Initially the patient experienced a loss in his business. As a result of this worry, he had depressed mood, low appetite and disturbed sleep reported after having the nightmares. In order to determine the patient's locus of control and degree of felt stress, the Rotter's Locus of Control Scale (Rotter, 1966), the Perceived Stress Scale (Cohen and Williamson, 1988), and the HADS (Zigmond and Snaith, 1983) were used. It was found that the patient had external locus of control. The score in depression came as 9 indicating mild depression and score in anxiety was 17 indicating presence of significant anxiety. In Perceived stress scale, the score was found to be 30 during baseline assessment which indicates severe level of perceived stress. Upon administration of Type A/B Behavior Pattern scale, it was found that the index patient had Type A (anxious) personality.

Hence, a cognitive behavioral therapeutic module addressing the anxiety, depression, stress and locus of control was planned to be done on the index patient. At first, the treatment focused on stress reduction using MBSR. Thereby patient's muscle tension was assessed using Electromyographic Biofeedback and he was trained to relax the muscles using mindfulness on respiratory biofeedback training so that he learns to self-induce relaxation by controlling his breathing rate. Once the patient was able to relax for a certain time period, the focus was kept on building his sense of autonomy in order to reduce the perceived loss of control. This included behavior activation whereby the patient engaged in doing household chores as well as finishing small targets daily of his business work within the designated time period. The family members also provided him with verbal reinforcement, on completion of such tasks. Once the patient did this for a

few weeks, it helped him build improved interpersonal relationship with his family members thereby improving his interpersonal social support network. At this point, the therapist initiated the discussion of the contents of his dreams with the patient. It was found that he had an unconscious need for aggression towards himself and an anxiety about his failing business. He also believed that he will not be able to take adequate care of his family members because of which he had extreme guilt which was expressed in his dreams in the form of his family members dying, as a form of passive aggression expressed in the dreams towards them. Hence, this was brought for discussion in session whereby the therapist focused on building self-efficacy in the patient with respect to adequately managing business, finances. In order to lessen the perceived family burden, his interpersonal relationship with wife was discussed focusing upon how responsibilities could be divided between the two as and when needed so that the patient does not feel overwhelmed and overburdened. Once the patient started practicing these at home, thereafter he reported having improved self-efficacy and decreased level of depression due to gaining efficiency in his work and better control upon his environment. Once he was able to understand that the choices and decisions of his life are within his control, it reduced his anxiety which was earlier raised due to perceived sense of loss of control. The patient also had difficulty in communicating his emotions to other people around him as he felt that others do not consider him to be worthy enough as an individual due to his poor productivity at work. Hence, Cognitive behavior therapy (CBT) was done on him to in order to target the cognitive errors which connected his sense of self-worth to his productivity. As a result, his negative affectivity also improved thereby decreasing the level of depression. After an intervention of about six months, it was found that the

levels of anxiety, depression and perceived stress in the patient have decreased to sub-clinical levels. The depression score came down to be 5 and anxiety score came down to be 9 which indicate that the anxiety has decreased from clinical level to borderline level. However, after follow-up of about one year, his scores in anxiety further came down to be 4 and depression score was at 3 which indicate that he did not have clinically significant depression and anxiety post follow-up. The locus of control was found to be more towards internal during follow-up as his sense of control over his surroundings improved due to improved coping. His frequency of nightmares also decreased to around twice in a month post intervention and after a one year follow-up, he reported that the nightmares have stopped occurring. It was also observed that he had a decrease in cardiac event as he reported having pain in the chest in around once in six months that is having it at a much lesser frequency than before. Hence, it could be said that the CBT Module has been an effective treatment method for this patient in reducing his nightmares related cardiac events.

### **Discussion**

The primary function of dreams is to facilitate the emotional processing of one's thoughts so that it can decrease the emotional reactivity in the patient. Dreams also help to articulate and blend the thoughts and anxieties together in a way that helps one to solve problems. The amygdala and hippocampus are areas of the brain that have been shown to have increased activity when we dream. However, dreaming facilitates creative problem-solving which can be concluded from the said effect on how people experience such brain events when sleeping which eventually facilitates the scientific understanding of how such events are experienced. The psychoanalysis theorizes that dreams represent cross-sectional images of a person's mental phenomenon

and anxieties relating to this such that the contents of dreams would directly relate to everyday activities, concerns, and interests of an individual. It becomes clear that dreams may not have clear meanings but rather be neural signals with subjective interpretation when the synthesis hypothesis is activated. This hypothesis was put forth by J. Alan Hobson in his Activation-synthesis hypothesis (Hobson, 2010), which postulates that dreams occur when random neurons in the brain fire during REM sleep and the mind tries to make sense of this firing. In the present case, it can be therefore said that dreams helped in the processing of emotions in the present case, and in that sense, it became a 'safe space' for the mind to resolve emotions and traumas that remain unresolved for him.

From a psychodynamic perspective, it may be conceptualized that the index patient harbored worries about his business failing and his wife and son suffering. Due to the Type A personality of the patient, he often remained anxious about daily life matters like interactions with colleagues and wife and also had minor conflicts with his children at times. He reported that his business started incurring minor losses since last two years. However he did not have significant worry over the matter except for the instances of nightmare in his dreams. This could indicate that the index patient repressed such worries as his it was beyond his ego strength to tolerate the anxiety. Due to repression, such thoughts which did not get processed adequately earlier, resulted in nightmares during sleep. Hence the imageries that he experienced appeared only in the dreams as nightmares because the process of repression prohibited it from appearing in the patient's conscious thought process during the wakeful state. Hence, in therapy, his dreams about the business were brought in for discussion in the sessions. The dream's thematic material is unique to the dreamer.

Therefore the dreamer's emotive reactions to the dreams are comparable to their affective reactions in real life. The patient ought to make an effort to summarize the dream and make a sense of its significance in his life (Carona and Fonseca, 2022). Thus, when it comes to reducing the frequency of nightmares, nightmare-focused cognitive behavioral therapy (such as exposure or systematic desensitization, cognitive restructuring, or imagery rehearsal therapy) has shown to be effective in the index patient. In the present case, the index patient's dreams was brought into discussion and analyzed in session in order to understand the sources of his anxiety and the same was intervened upon in the session which eventually brought about a reduction in nightmares and frequency of cardiac events. A web-based cognitive behavioral therapy for insomnia (wCBT-I) demonstrated feasibility and resulted in a clinically meaningful improvement in the severity of insomnia in an older patient group with a high prevalence of CAD (Javaheri, Reid, Drerup et. al, 2019), a finding similar to what we have found in our current study. Hyperlipidemia, a risk factor for coronary artery disease, may be linked to recurrent nightmares in older persons (Lee, Park, Kim et. al., 2024). CAD is linked to the frequency and intensity of nightmares experienced by veterans. According to study results, nightmares can be a stand-alone risk factor for Coronary artery disease (Campbell, Taylor, Augustine et. al., 2023). Hence it can be seen from such research findings that nightmares is so intricately related to the onset and maintenance of cardiac events and therefore intervening upon the index patient's nightmares proved to be crucial and effective in reducing his frequency of cardiac events.

It was found that whenever the patient actively discussed about the matters of business for a prolonged duration, he experienced pain in the chest. Hence, he

repressed such emotions into the unconscious. However, studies show that masking and repressing emotions may trigger physiological responses to stress. Stress arises from the social condemnation and punishment of open expression of emotions that lead to repression which is itself intimidating and stressful. According to Cote (2005), persistent repression causes stress in people, which can have negative impacts on an individual's productivity. These effects include elevated heart rate, anxiety, low commitment, and other issues. That might be the reason why when the index patient had heightened stress due to Type A personality characteristics, he was unproductive at his work, thereby serving to increase his stress further. Over time, the level of stress increased so much that it exceeds the capacity of his ego-strength. Then, his conscious mind or the ego represses it into the unconscious, hence the index patient was not aware of feeling of stress actively in his daily life but nevertheless dreamt about the same. When under stress, he experienced chest pain because of the condition's impact on his blood flow and the body's reaction to stress. Additionally, stress used to trigger the release of hormones like cortisol and adrenaline, which raised blood pressure and heart rate and make breathing harder for him. Thus his chest pain might have also result from these impacts. Stress can also release cortisol hormone in the blood which might be responsible for increased blood pressure and rupture in plaque, forming blood clots and therefore blocking adequate and timely blood flow and thereby might be causing a heart attack in individuals with Coronary artery disease. However this psychotherapeutic module proved to be effective for the index patient as it helped to bring out the repressed material on the surface of consciousness and deal with it in session in order to bring about decreased stress, anxiety and therefore improve coping and mood state, which in turn

might have also positively impacted or reduced the frequency of cardiac events in the index patients.

#### Limitations of the study

It is however imperative to state that the current study has certain limitations. Firstly since it is a single-case experimental design, therefore its findings cannot be generalized on a broader population which includes people belonging to a different state, country or socioeconomic background. Secondly, even though CBT was effective for treating Nightmares, it cannot be said for certain whether CBT will work for other kinds of sleep disorders as well in patients having coronary artery disease. However, despite the limitations, the study shows how sleep dysfunction can be related to and can further complicate the picture of coronary artery disease and it provides a good framework of demonstration for treating sleep disorders using psychodynamic approach in one such cardiac patient.

#### Conclusion

The present study shows that the incidence and severity of coronary artery disease has been significantly impacted by dreams in the index patient. Nightmares are an expression of the repressed distress in the patient, which in turn further led to increased stress thereby increasing the level of cortisol in the blood, serving to cause and maintain coronary artery disease. Therefore, understanding the content of dreams in nightmares and intervening upon them served to decrease the level of anxiety in the patient, thereby reducing his stress level, which might have led to further reduction in frequency and intensity of the cardiac event in the index patient. The findings of the present study may therefore be used in patients having CAD alone as well in patients who have nightmares and other such psychological disorders which exacerbated the symptoms of Coronary Artery Disease.

#### References

- Campbell, A. A., Taylor, K. A., Augustine, A. V., Sherwood, A., Wu, J. Q., Beckham, J. C., Hoerle, J. M., VA Mid-Atlantic MIRECC, Ulmer, C. U (2023) Nightmares: an independent risk factor for cardiovascular disease?, *Sleep*, 46(6). <https://doi.org/10.1093/sleep/zsad089>
- Carona, C., & Fonseca, A. (2022). A clinical approach to nightmares and bad dreams in cognitive-behavioural therapy. *British Journal of Psychiatry Advances*, 28(3), 195–197. doi:10.1192/bja.2021.77
- Cohen, S., & Williamson, G. (1988). Perceived stress in a probability sample of the United States. In S. Spacapan & S. Oskamp (Eds.), *The social psychology of health: Claremont Symposium on applied social psychology*. Newbury Park, CA: Sage.
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of health and social behavior*, 24(4), 385–396.
- Cote S. (2005) A social interaction model of the effects of emotion regulation on work strain. , *Academy of management review* 30: 509-530.
- Dhar, U., & Jain, M. (2001). Manual for Type A/B Behavior Pattern Scale. Ankur Psychological Agency, Lucknow
- Hobson, J. Allan (2010). "REM sleep and dreaming: Towards a theory of protoconsciousness". *Nature Reviews Neuroscience*. 10 (11): 803-13 . [doi: 10.1038/nrn2716](https://doi.org/10.1038/nrn2716). PMID 19794431. S2CID 205505278
- Javaheri, S., Reid, M., Drerup, M., Mehra, R., & Redline, S. (2019). Reducing Coronary Heart Disease Risk Through Treatment of Insomnia Using Web-Based Cognitive Behavioral Therapy for Insomnia: A Methodological Approach. *Behavioral Sleep Medicine*, 18(3), 334–344. <https://doi.org/10.1080/15402002.2019.1584896>
- Kumar, P. N. S., & Thomas, B. (2007). Family intervention therapy in alcohol dependence syndrome: One-year follow-up

- study. *Indian journal of psychiatry*, 49(3), 200–204. <https://doi.org/10.4103/0019-5545.37322>
- Lee, Y., Park, D., Kim, S., Shin, C., & Suh, S. (2024). Association of nightmares with cardio-cerebrovascular disease, hypertension and hyperlipidemia in older adults: A population-based cross-sectional study. *Journal of psychosomatic research*, 182, 111669. <https://doi.org/10.1016/j.jpsychores.2024.111669>
- Li, Y. N., Buys, N., Ferguson, S., Li, Z. J., & Sun, J. (2021). Effectiveness of cognitive behavioral therapy-based interventions on health outcomes in patients with coronary heart disease: A meta-analysis. *World journal of psychiatry*, 11(11), 1147–1166. <https://doi.org/10.5498/wjp.v11.i11.1147>
- Ressler K. J. (2010). Amygdala activity, fear, and anxiety: modulation by stress. *Biological psychiatry*, 67(12), 1117–1119. <https://doi.org/10.1016/j.biopsych.2010.04.027>
- Rotter, J. B.: (1966). Generalized expectancies for internal versus external control of reinforcement. *Psychological monographs: General and applied*, 80(1):1.
- Vincent N, Dirkse D, Giannouli E, McQuarrie A. (2023) Transdiagnostic cognitive behavioral therapy for nightmares and parasomnias. *Journal of Clinical Sleep Medicine*.19(3):499–509.
- Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta psychiatrica Scandinavica*, 67(6), 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x>

**Usri Sengupta**, M. Phil in Clinical Psychology from RINPAS, currently PhD Scholar at the Department of Clinical Psychology, RINPAS.Ranchi and Clinical Psychologist, Department of Psychiatry, ICMR-BMHRC

**Amool Ranjan Singh**, PhD in Clinical Psychology. Professor and Head, Department of Clinical Psychology, Ranchi Institute of Neuro-Psychiatry and Allied Sciences (RINPAS), Ranchi

**Hemant Narayan Ray**, MD in Cardiology, Professor and Head, Department of Cardiology, Rajendra Institute of Medical Sciences (RIMS), Ranchi